

## Ciprofloxacin -Tablets

Strength	Composition	Therapeutic Segment	Class
250 mg	Ciprofloxacin HCl BP Eq. to Ciprofloxacin 250 mg	Antibiotic	Fluoroquinolone
500 mg	Ciprofloxacin HCl BP Eq. to Ciprofloxacin 500 mg		
<b>Indications:</b> UTI, Acute Uncomplicated Cystitis in females, Chronic Bacterial Prostatitis, LRTI, Acute Sinusitis, SSTI, Bone and Joint Infections, Complicated Intra-Abdominal Infections, Infectious Diarrhea, Typhoid Fever (Enteric Fever)			
<b>Side-effects:</b> Nausea , diarrhea, liver function tests abnormal, vomiting and rash			
<b>Precautions:</b> Epilepsy, CNS disorder, renal or hepatic dysfunction, G6PD deficiency, good hydration required myasthenia gravis, may impair ability to drive or operate machinery			
<b>Contra-indications:</b> Children < 12 years, Pregnancy, lactation, avoid exposure to sunlight or sun lamps			
<b>Overdose:</b> stomach should be emptied by inducing vomiting or by gastric lavage. supportive treatment, Adequate hydration			
<b>How to take:</b> Do not take with dairy products such as milk . Take this medication for as many days as it has been prescribed			