

## Co-Trimoxazole Tablets

Strength	Composition	Therapeutic Segment	Class
480 mg	Trimethoprim BP 80 mg Sulphamethoxazole BP 400 mg	Antibiotic	Dihydrofolate reductase inhibitor/ Sulfonamide antibiotic
960 mg	Trimethoprim BP 160 mg Sulphamethoxazole BP 800 mg		
<p><b>Indications:</b> Urinary Tract Infections, Acute Otitis Media, Acute Exacerbations of Chronic Bronchitis in Adults, Travelers' Diarrhea in Adults, Shigellosis, Pneumocystis Carinii Pneumonia</p>			
<p><b>Side-effects:</b> Gastrointestinal disturbances (nausea, vomiting, anorexia) and allergic skin reactions (such as rash and urticaria).</p>			
<p><b>Precautions:</b> G6PD deficiency, folate deficiency, hepatic and renal impairment, elderly, porphyria, thyroid dysfunction, maintain adequate hydration</p>			
<p><b>Contra-indications:</b> Hypersensitivity, severe hepatic or renal insufficiency, infants &lt; 2 months, Pregnancy, Lactation</p>			
<p><b>Overdose:</b> Gastric lavage or emesis, intravenous fluids, Acidification of the urine, blood counts and blood chemistries</p>			
<p><b>How to take:</b> Take co-trimoxazole with food.</p>			